# *ARDEA STELLARIS CRISTATA AMERICANA.*

## *The creſted Bittern*

WEIGHS a Pound and half. The Bill is black and ſtrong. The Eyes very large and prominent, with red *Irides*. The Skin encompaſſing the Eyes is green. The Crown of the Head, from the Baſis of the Bill, of a pale yellow, terminating in a Peak; from which hang three or four long white Feathers, the longeſt of which is ſix Inches; which they erect, when irritated. From the Angle of the Mouth runs a broad white Liſt. The Reſt of the Head is of a blueiſh black. The Neck, Breaſt and Belly duſky blue. The Back is ſtriped with black Streaks, with a Mixture of white. From the Upper-part of the Back ſhoot many long narrow Feathers, extending beyond the Tail; ſome of which are Seven Inches long. The large Feathers of the Wing are brown, with a Tincture of blue. The Legs and Feet are yellow. Theſe Birds are ſeen in *Carolina* in the rainy Seaſons; but in the *Bahama* Iſlands, they breed in Buſhes growing among the Rocks in prodigious Numbers, and are of great Uſe to the Inhabitants there; who, while theſe Birds are young, and before they can fly, employ themſelves in taking them, for the Delicacy of their Food. They are, in ſome of theſe rocky Iſlands, ſo numerous, that in a few Hours, two Men will load one of their *Calapatches* or little Boats, taking them pearching from off the Rocks and Buſhes; they making no Attempt to eſcape, tho' almoſt full grown. They are called, by the *Bahamians*, Crab-catchers, Crabs being what they moſtly ſubſiſt on; yet they are well-taſted, and free from any rank or fiſhy Savour.

## Butor huppé.

*IL pèſe une livre & demie. Son bec eſt noir & fort. Ses yeux ſont fort grands, & protuberans, avec des* Iris *rouges. La peau qui entoure ſes yeux, eſt verte. Le deſſus de ſa tête, depuis la baſe de ſon bec, eſt d'un jaune pâle, qui ſe termine dans une pointe, d'où partent trois ou quatre longues plumes blanches, dont la plus longue eſt de ſix pouces. Lorſque l'oiſeau eſt en colere, il dreſſe ces plumes. Il part du coin de ſon bec une large raye blanche. Le reſte de ſa tête eſt d'un noir tirant ſur le bleu. Son cou, ſa poitrine & ſon ventre ſont d'un bleu obſcur. Son dos eſt rayé de noir & de blanc. Il naît pluſieurs longues plumes étroites à la partie ſuperieure de ſon dos, qui s'étendent au delà de ſa queue. Quelques unes de ces plumes ont ſept pouces de long. Les grandes plumes de l'aîle ſont brunes, avec une nuance de bleu. Ses jambes & ſes pieds ſont jaunes. On voit ces oiſeaux à la* Caroline *dans la ſaiſon des pluyes: Mais aux Iſles* Bahama, *ils ſont en très grand nombre, & font leurs petits dans des buiſſons qui viennent parmi les rochers. Les gens du Païs ſçavent bien en faire leur profit, car tandis que ces oiſeaux ſont jeunes, & avant qu'ils puiſſent voler, ils s'amuſent à les prendre pour s'en regaler. Ils ſont en ſi grand nombre dans quelques unes de ces Iſles pleines de rochers, qu'en peu d'heures, deux hommes en prendront aſſez pour charger un de leurs petits bateaus. Ils ſe laiſſent prendre de deſſus les rocs & les buiſſons ſur leſquels ils ſont perchez, ſans faire mine de s'enfuir, quoique déja grands. Les* Bahamiens *les appellent Prenneurs de Cancres, ce coquillage étant preſque leur ſeule nourriture; cependant ils ſont d'un très bon goût, & ne ſentent en aucune maniére le marécage.*

## *Lobelia fruteſcens, Portulacæ folio.* Plum. Nov. Gen. p. 21.

THIS Plant grows uſually to the Height of five or ſix Feet. The Leaves are, in Thickneſs and Form, not unlike Purſlain. At the End of a Stalk, growing from the Joint of a Leaf, there are ſet three or four monopetalous white Flowers, divided into five pointed Sections, with a wreathed *Stamen* hanging out. The Flower here exhibited is ſlit down to the Baſis and laid flat open. The Flowers are ſucceeded by globular Berries, of the Size of black Bullace, containing a Stone, covered with a ſmooth black Skin. Theſe Plants grow on the rocky Shores of many of the *Bahama* Iſlands.

*CETTE plante croît ordinairement à la hauteur de cinq ou ſix pieds. La figure & l'épaiſſeur de ſes feuilles reſſemblent fort au Pourpier. Au bout d'une tige, qui naît de l'aiſſele d'une feuille, ſont attachées trois ou quatre fleurs blanches monopétales, diviſées en cinq ſections pointuës, avec une étamine torſe qui pend en dehors. Cette fleur paroît d'une maniére particuliére. Il ſemble qu'elle ait été tubuleuſe, mais fenduë juſqu'à la baſe, ouverte & applatie. Ces fleurs ſont ſuivies par des bayes rondes, de la groſſeur des prunelles noires. Elles contiennent un noyau couvert d'une peau noir & unie. Ces plantes croiſſent ſur les rochers qui ſont ſur les bords des Iſles* Bahama.